

RESEARCH REPORT

EFFECTS OF MYOFASCIAL RELEASE AND VAPOCOOLANT SPRAY WITH STRETCH TECHNIQUE ON UPPER TRAPEZIUS TRIGGER POINTS

ABSTRACT

BACKGROUND

Myofascial pain syndrome is a common source of pain in people with bad postures. Mostly trigger points (TrP) of muscles become chronic and cause restricted ROM. Upper trapezius TrP are common in people of all ages especially young and middle aged adults. Physical therapy has been used to release these TrP.

OBJECTIVES

To determine the effects of Myofascial release and spray with stretch technique on upper trapezius trigger points

METHODOLOGY

30 individuals with history of neck pain were taken through non-probability purposive sampling technique. They are randomly allocated to intervention and control group. The intervention group received Myofascial release technique with stretching exercises and the control group received spray with stretch technique. The techniques were applied for three consecutive days. A pre-assessment and post-assessment of ROM and pain scores on VAS were taken.

RESULT

This study shows that the ranges of cervical side flexion of both sides were increased in intervention and control groups.

CONCLUSION

Myofascial release have greater effects in releasing upper trapezius TrP with significant reduction in neck pain and improved neck movements as compared to spray and stretch technique and there would be minimal chances of early recurrence of TrP after about 3 days of receiving it.

KEYWORDS

Trigger Points, Myofascial Release, Spray and Stretch Technique, Upper Trapezius, Neck Pain, Range of Motion, VAS

Fareeda Shaheen Shah

Consultant Physiotherapist
Department of Physiotherapy
Sindh Government Hospital,
Saudabad
fareeda.shah08@gmail.com

Ali Farhad

Senior Lecturer
Ziauddin College of Physical Therapy
Ziauddin University
alifarhadtharwani@gmail.com

Syeda Najaf Zehra

DPT Student
Ziauddin College of Physical Therapy
Ziauddin University

Zoya Jabeen

DPT Student
Ziauddin College of Physical Therapy
Ziauddin University

Tayyab Tahir

DPT Student
Ziauddin College of Physical Therapy
Ziauddin University

[Shah FS, Farhad A, Zehra SN, Jabeen Z, Tahir T. Effects of Myofascial Release and Vapocoolant Spray with Stretch Technique on Upper Trapezius Trigger Points. Pak. j. rehabil. 2016;5(2):43-48]

INTRODUCTION

Myofascial trigger point is a hyperirritable spot that is felt as a nodule in a taut band of skeletal muscle¹. Manual pressure is used to locate the site of trigger points (TrP)¹. These trigger points cause Myofascial pain syndrome and have sensory, motor and autonomic dysfunction¹. Motor symptoms include decrease muscle strength with stiffness and decrease range of motion while sensory symptoms manifest as local tenderness, referred pain with central and peripheral sensitization¹. Central sensitization refers to increased excitability of central neurons whereas peripheral sensitization describes the reduction in threshold and increase nociceptors responsiveness¹. These sensitizations lead to allodynia and hyperalgesia¹. Autonomic symptoms include altered vasoconstriction and vasodilatation, coldness, increased sweating and pilomotor response with hypersecretion².

The potential causes of trigger points include overuse activities, trauma or sudden acute overloading of muscle, muscle deconditioning, postural imbalances associated with poor ergonomics, asymmetrical loading while lifting heavy objects³.

Myofascial trigger points can be active that produce classic pattern of pain and can be latent that are asymptomatic⁴. Further types include primary or secondary trigger points in which the primary is due to acute or chronic overloading while secondary develops as a result of primary trigger point and satellite or associated trigger points that are formed in a reference zone of other trigger points⁴. The radiating pain of trigger points is generally recognizable and the body area to which the pain radiates is known as its reference zone⁴. The radiating pain is dull and aching in nature associated with or without discomfort. Palpation is done to examine the tenderness as latent trigger points cause pain on palpation that is otherwise asymptomatic⁴.

Poor head posture leads to hyperactivity in upper trapezius that causes trigger point formation in chronic stages⁴. The associated trigger points can be formed in the contralateral trapezius and ipsilateral levator scapulae, rhomboids and supraspinatus muscles⁴. Moreover, the satellite trigger points are more likely to develop in temporalis and occipitals muscle⁴. Trapezius has three parts that are pars descendens, parstransversa, pars ascendens⁵. Pars descendens is the upper part of trapezius that originates from the occipital protuberance, nuchal ligament, spinous processes of 7th cervical and all thoracic vertebrae and are inserted on the lateral third of the clavicle and acromion and spine of the scapula⁵. The fibers of the upper part elevate the acromial end of the clavicle, extend the head and stabilize the scapula⁵. There are seven motor points

in the trapezius muscle among which four are the motor points that are clinically associated with trigger point₁ (TrP₁)⁶. Trigger point₂ (TrP₂) is located in the mid portion of anterior border of upper trapezius⁴. The involved fibers are vertical that attached to the clavicle⁴. The referred pain goes from the posterolateral aspect of neck to the mastoid process⁴. This referred pattern is associated with 'tension headache'⁴. The pain is rarely referred to the lateral side of head, temple, back of orbit and jaw angle⁴. TrP₂ is inferior and lateral to the TrP₁ in the mid horizontal fibers⁴. The resultant pain is referred behind the ear and posterior to the zone of TrP⁴.

Various techniques are used to treat the trigger points such as ice massage, ice sweeps, spray and stretch, sustained deep pressure, electrical stimulation, low-power laser and ultrasound⁷. Application of cold alleviates the pain by decreasing nerve conduction velocity (the speed by which a nerve signal travels)⁸. Also it acts as a counterirritant and desensitizes the nociceptors according to pain gate theory; another effect of decrease in pain is due to the release of endorphins from the descending pathways⁸. Myofascial release refers to a manual massage technique that is applied to release fascia, skin, soft tissues and bone from adherence in order to relieve pain and to improve range of motion⁹. Fascia is a connective tissue that lines the musculoskeletal system and organs as well⁹.

Myofascial release encompasses different manual techniques among which sustained manual pressure technique has found to be most effective treatment for releasing trigger points¹⁰. A study hypothesized that the central nervous system is involved in the pathogenesis of trigger points¹¹. This central mechanism explains prolonged plateau depolarization of α -motor neuron that results in continuous discharge from the neurons¹¹. Myofascial pain syndrome is a musculoskeletal syndrome that is associated with trigger points in the muscles¹². These trigger points are palpable, focal and tender¹². On palpation they produce twitch response with or without referral of pain¹².

Myofascial release is a manual technique that implies application of gentle manual pressure to release the fascia adhesions¹³. The technique is applied without any ointment into the direction of adherence¹³. The pressure is sustained for about 90 to 120 seconds minimally¹³. Myofascial release technique restores muscular function with postural alignment and promotes relaxation⁹. Basically this technique releases the muscles that are held abnormally with a tight fascia⁹.

Researchers supported the idea of releasing trigger points with vapocoolant sprays¹⁴. The evidences have explained that the application of cold by the spray inhibits the central nervous system responses including pain signals along with reflex motor and

autonomic effects¹⁴. As the pain is alleviated the muscles become relaxed which can be then stretched and lengthened¹⁴.

This spray and stretch technique is effective, quick and least painful for the treatment of a single muscle myofascial pain¹⁵.

Vapocoolant spray provides tactile stimulation in the reference zone of trigger points that blocks the reflex spasm¹⁵. Thus the pain subsides and the muscle is stretched passively¹⁵. The main purpose of this study was to see the effects of vapocoolant spray with stretching on myofascial trigger points.

METHODS

Study Design

Randomized control trial (RCT)

Population and Study Sample

30 participants who aged between 18 to 40 years were recruited in the study. A block randomization non-probability purposive sampling technique was used according to which 15 participants were allocated to intervention group and 15 to control group.

Sources of Data

Data was collected from an OPD setting of an orthopedic clinic.

Collection of Data

Data was collected through questionnaires and assessment forms. On the basis of questionnaire the participants were recruited into the study. After recruitment a pre-assessment was taken including VAS score and range of motion for cervical lateral flexion. Pain was elicited by palpating the trigger points and the participants were asked to mark the pain intensity on a VAS scale. Participants who notified bilateral pain on palpation were assessed for range of motion of cervical lateral flexion on both sides that indicated bilateral involvement with trigger point formation.

Procedure

Subjects were equally allocated to two groups A and B. Group A received Myofascial release technique with neck stretching exercise for upper trapezius whereas group B received spray and stretch technique with neck stretching exercises. The treatment was applied for three days consecutively and in each session technique was applied thrice. Post-assessment was done after third session.

Group A:

• Myofascial Release

Subjects were instructed to sit in a comfortable position with back straight and supported on the armless chair and feet placed on the floor. Sustained manual pressure was applied for 3 times

per session. First the trigger points were palpated in the upper fibers of trapezius by thumb moving in a circular direction deeply in different points of upper trapezius, when found deep sustained pressure on the trigger points was applied with thumb and maintained for 60 seconds and then was released in vertical direction.

• Stretching of upper trapezius muscle

Therapist laterally flexed the neck followed by forward flexion on opposite side to place stretch on the involved upper fibers of trapezius muscle.

Group B: Spray and stretch

Subjects were instructed to sit in a comfortable position with back straight and support on the armless chair and feet placed on the floor. Then the therapist stretched the upper fibers of trapezius muscle for maximum lengthening with the Myofascial trigger point and pressed the patient's head in forward direction to raise the occiput. At the same time the vapocoolant spray was applied in parallel sweeps from the acromion to the mastoid process.

Data analysis strategies

A Statistical package for social sciences (SPSS 20) version has been used for the analysis of statistical data.

Inclusion criteria

Individual with history of neck pain or tension headache for at least 3 months and were willing to participate in the study.

Exclusion criteria:

Individuals with who have one of the following conditions were excluded from the study.

- open wounds
- Fever
- Skin infections
- Impaired circulation
- Disc herniation
- Recent spine surgery
- History of cervical fracture

RESULTS

This study analyzed the comparative effects between two techniques that are Myofascial release with sustained pressure and spray and stretch techniques in the treatment of Myofascial pain syndrome caused by upper trapezius trigger point. Stretching exercises of neck are incorporate with both techniques. Participants were selected on the basis of questionnaires and assessment (i.e. VAS score and range of motion of cervical side flexion). Sample of 30 participants were included in the study with mean age of 18 to 40 years. It was seen that out of 30 randomly selected patients, 21 (70%) were female and 9 (30%) were male.

Table 1

P<0.05 is considered as significant in paired t-test (Table 1)

Myofascial Release	Preassessment	Postassessment	
	Mean ± std. Deviation	Mean ± std. Deviation	p-value
Pair-1 VAS	6.07±1.710	1.93±1.033	<0.01*
Pair-2 right cervical side flexion	36.5333±7.33744	46.3333±8.26928	<0.01*
Pair-3 left cervical side flexion	36.467±7.4942	44.3333±7.65942	<0.01*

Table 2

p<0.05 is considered as significant in paired t-test

Spray and stretch technique	Preassessment	Postassessment	
	Mean ± std. Deviation	Mean ± std. Deviation	p-value
Pair-1 VAS	5.00±1.604	2.93±1.335	<0.01*
Pair-2 right cervical side flexion	38.000±6.23584	44.8667±5.68038	<0.01*
Pair-3 left cervical side flexion	38.9333±9.92448	43.2667±7.14609	0.06

Patients were equally divided into two groups i.e. 15 patients per group. Group A was treated by myofascial release with exercises while group B was treated by spray and stretch technique with exercises. Pre and post range of motion of cervical side flexion were noted of the respective treatment given. The data was analyzed from the SPSS statistical analysis program on the basis of mean and standard deviation by using t-table.



It is transpired from the descriptive analysis that mean range of Myofascial release with exercises pre and post cervical side flexion of right side are 36.53 and 46.33, which shows an increase in range of 9.8 in right cervical side flexion while the mean range of myofascial release with exercises pre and post cervical side flexion of left side are 36.46 and 44.33 which shows an increase in range of 7.87 in left cervical side flexion.

At the same time, mean range of spray and stretch technique with exercises pre and post cervical side flexion of right side are 38.00 and 44.86 which shows an increase in range of 6.86 in right cervical side flexion while the mean range of spray and stretch technique with exercises pre and post cervical side flexion of left side are 38.93 and 43.26 which shows an increase in range of 4.33 in left cervical side flexion.

Pain was also calculated via Visual Analogue Scale (VAS) in candidates of both groups i.e. A and B. Group A, calculated pain was 6.07 before treatment and 1.93 after treatment. While group B, pretreatment pain was 5.00 and post treatment pain becomes 2.93.

By above results, it was known that both these techniques are effective in relieving pain and restoring range of motion in participants with myofascial pain syndrome caused by upper trapezius trigger point with p-value 0.00 by applying paired sample test.

In comparison, it is evident that results of myofascial release with exercises in terms of pain reduction and increase in range of motion are more significant as compared to the results of treatment with spray and stretch technique with exercises.

Myofascial Release with sustained Pressure**Spray and Stretch Technique**

DISCUSSION

Myofascial pain syndrome is a major health issue that affects the individuals by reducing mobility with disabling pain. It therefore has impact on sense of well-being¹⁶. This syndrome comprises the painful taut bands of muscle that contain palpable, hyper-sensitive nodules known as myofascial trigger points¹⁶. It has been reported that individuals with trigger points not only show a reduced functional status but also suffer from mood changes and have an indirect effect on quality of life¹⁶.

Individuals who have trigger points report local persistent pain with decreased range of motion¹⁷. Mostly the trigger point formation occurs in postural muscles¹⁷. Postural muscles of the neck are the most frequently affected ones including trapezius especially¹⁷. It has been suggested that active trigger points in the upper trapezius muscle were associated with chronic tension type headache. It was also reported that individuals with active trigger points had been suffering with more severe type of tension headache as compared to those having latent trigger points¹⁸. On Histological examination of muscle biopsies from myofascial trigger points reveals that hyper contracture is formed that is associated with sustained sarcoplasmic reticulum calcium release due to intense neural activation and action potential generation¹⁶. Sustained contraction with increased metabolic stresses and reduced blood flow causes secondary changes in the muscle and leads to persistent trigger points¹⁶.

The proposed mechanism behind myofascial release with sustained pressure is basically the neuromuscular relaxation. This relaxation is achieved by the stimulation of Golgi tendon organs which alter the mechanical tension of the muscle with resultant reduction in neuromuscular sensitivity to mechanical pressure and stretch¹⁸.

In a 1998 study, Simons concluded that spray and stretch technique is the most effective noninvasive technique in deactivating trigger points¹⁹. A temperature reduction of 2 to 5 degrees is enough to cool the tissues and the resultant beneficial effects include analgesic, vasomotor, anti-inflammatory and muscle relaxation effects^{19,20}. According to Travell and Simons, vapocoolants have a cooling effect on the area being affected. This causes the transmission of sensory impulses to the spinal cord that result in an inhibitory effect on local pain. The same inhibitory effects actually break the pain spasm pain cycle due to which the muscle ultimately becomes relax and the stretching further aids in elongation of the muscle¹⁸. In Spray and Stretch technique, the affected muscle is passively stretched^{21,22} and dichlorodifluoromethane-trichloromonofluoromethane (Fluori- Methane) or ethyl chloride spray is applied simultaneously on the skin.¹⁸

The mean range of Myofascial release with exercises pre and post cervical side flexion of right side are 36.53 and 46.33, which shows an increase in range of 9.8 in right cervical side flexion. And the mean range of myofascial release with exercises pre and post cervical side flexion of left side are 36.46 and 44.33 which shows an increase in range of 7.87 in left cervical side flexion.

At the same time, mean range of spray and stretch technique with exercises pre and post cervical side flexion of right side are 38.00 and 44.86 which shows an increase in range of 6.86 in right cervical side flexion. And the mean range of spray and stretch technique with exercises pre and post cervical side flexion of left side are 38.93 and 43.26 which shows an increase in range of 4.33 in left cervical side flexion.

Pain was also calculated via Visual Analogue Scale (VAS)²³⁻²⁵ in candidates of both groups i.e. A and B. Group A, calculated pain was 6.07 before treatment and 1.93 after treatment. While group B, pretreatment pain was 5.00 and post treatment pain becomes 2.93.

It was acknowledged that both these techniques are effective in relieving pain and restoring range of motion in participants with myofascial pain syndrome caused by upper trapezius trigger point with p-value 0.00 by applying paired sample test.

REFERENCES

- [1] Dommerholt J, Bron C, Franssen J. Myofascial Trigger Points. An Evidence-Informed Review. *J Man Manip Ther.* 2006;14(4):203-221
- [2] Benjamin PJ, Tappan FM. Tappans Handbook of Healing Massage Techniques. 5th Ed. Prentice Hall; 2009
- [3] Kisner C, Colby LA. Therapeutic Exercise Foundations and Techniques. 6th Ed. Philadelphia: FA Davis Company; 2012
- [4] Simons DG Travel JG, Simons LS. Myofascial Pain and Dysfunction the Trigger Point Manual Volume 1.Upper Half of Body. 2nd Ed. Philadelphia: Williams & Wilkins; 1999
- [5] Physio-Pedia.Com. Lectures and Presentation-Physiopedia, Universal Access to Physiotherapy Knowledge. [Internet]. 2015[Cited 4 September 2015]
- [6] Akamatsu FE, Ayres BR, Saleh SO, Hojaij F, Andraid M, Hsing WT et al. Trigger Points: An Anatomical Substratum. *BioMed Res Int.* 2015;2015:623287
- [7] Michlovitz SL, Bellew JW, Nolan Jr TP. Modalities for Therapeutic Intervention. 5th Ed. Philadelphia: F.A Davis Company; 2012
- [8] Algafly AA, George KP. The Effect of Cryotherapy on Nerve Conduction Velocity, Pain Threshold and Pain Tolerance. *Br J Sports Med.*

- 2007;41(6):365-369
- [9] Shah S, Bhalara A. Myofascial release. *Internat J Health Sci Res.* 2012;2(2):69-77
- [10] Fryer G, Hodgson L. The Effect of manual pressure release on myofascial trigger points in the upper trapezius muscle. *J Bodyw Mov Ther.* 2005;9(4):248-255
- [11] Hocking MJ. Exploring the central modulation hypothesis: do ancient memory mechanisms underlie the pathophysiology of trigger points? *Curr Pain Headache Rep.* 2013;17(7):347
- [12] Frante F, Bearn L, Rothrock R, King L. Evidence against trigger point injection technique for the treatment of cervicothoracic myofascial pain with botulinum toxin type A. *Anesthesiology.* 2005;103(2):377-383
- [13] Lebauer A, Brtalik R, Stowe K. The Effect of myofascial release on an adult with idiopathic scoliosis. *J Bodyw Mov Ther.* 2008;12:356-363
- [14] Kostopoulos D, Rizopoulos K. Effect of topical aerosol skin refrigerant (Spray and Stretch technique) on passive and active stretching. *J Bodyw Mov Ther.* 2008;12(2):96-104
- [15] Simons D. *Myofascial Pain Syndrome Due To Trigger Points.* Cleveland: Ohio; 1983
- [16] Jafri M. *Mechanisms of Myofascial Pain.* International Scholarly Research Notices. 2014;2014:1-16
- [17] Alvarez D, Rockwell P. triggers points: diagnosis and management. *American family physician.* 2002;65(4):653-660
- [18] Doraisamy MA, Anshul CPK, Gnanamuthu C. Chronic tension type headache and the impact of myofascial trigger point release in the short term relief of headache. *Glob JHealth Sci.* 2010;2(2):238-234
- [19] Spray And Stretch Revisited - The Little Secret With Trigger Points. Sep 2007. Available from: <https://www.bodybuilding.com/fun/stalely16.htm>
- [20] Trigger Point Therapy-Spray and Stretch Techniques. Dec 2016. [Internet]. Neil Asher Healthcare. Available from: <https://www.nielasher.com/blogs/video-blog/73449605-trigger-point-therapy-spray-and-stretch-techniques>
- [21] Williams JG, Laudner KG, McLoda T. The acute effects of two passive stretch maneuvers on pectoralis minor length and scapular kinematics among collegiate swimmers. *Int J Sports Phys Ther.* 2003;8(1):25-33
- [22] Ahmed H, Iqbal A, Anwar S, Alghadir A. Effect of modified hold-relax stretching and static stretching on hamstring muscle flexibility. *J Phys Ther Sci.* 2015;27(2):535-538
- [23] Haefeli M, Alfering A. Pain assessment. *Eur Spine J.* 2006;15(1):17-24
- [24] Baeyer CL, Bergstrom KJ, Brodwin MG, Brodwin SK. Invalid use of pain drawings in psychological screening of back pain patients. *Pain.* 1983;16:103-107
- [25] Bird SB, Dickson EW. Clinically significant changes in pain along the visual analog scale. *Ann Emerg Med.* 2001;38:639-643

