ANALYSIS OF FACTORS LEADING TOWARDS OBESITY AMONG SCHOOL - GOING CHILDREN (AGED 13 - 16 YEARS) OF KARACHI

ABSTRACT

OBJECTIVES

To analyze the factors leading to obesity in school children and to assess the role of Physical therapist in health education in schools

METHODS

It was a cross sectional study and factors leading towards obesity among school - going children aged 13 - 16 years were analyzed. A total of 300 students were selected from different schools of Karachi through Convenience sampling technique. Data entry was done on SPSS version 20 and frequencies and percentages were taken as descriptive statistics for categorical variables.

RESULTS

300 participants were included in the study where girls were 53.7% and boys were 46.3%. It was found that majority 40% of the children come under the category of normal BMI while 37% were overweight and obese. In addition, 55% of the children had a positive family history of obesity. It was also interesting to find out that 46% students were not participating in any physical activity and 45% schools didn't even had the PT period in their schedule. Furthermore, 91% students were consuming regularly.

CONCLUSIONS

Our study concluded that the major risk factor of obesity in 13-16 years children is lack of physical activity. However, usage of junk food is also increasing in this age group. Furthermore genetic predisposition cannot be ignored. It was further noticed that most of our children are having less sleep than recommended which itself has a very strong relation with weight gain.

Key Words

Physical Activity, Obesity, Children, School, Teenage, Overweight.

Muhammd Sarfraz

Assistant Professor
Ziauddin College of Physical Therapy
Ziauddin University
mohdprk23@hotmail.com

Anum Tariq

Lecturer
Ziauddin College of Physical Therapy
Ziauddin University
anumtariq.88@hotmail.com

Shahnaila Aziz

Physiotherapist
Aga Khan Social Welfare Board
Aga Khan Hospital & University
Preeti_shani@hotmail.com

Abida Nadeem

Physiotherapist
Al Fatah Medical Center
abidanadeem39@yahoo.com

[Sarfraz M, Tariq A, Aziz S, Nadeem A. Analysis of Factors Leading towards Obesity among School - Going Children (aged 13 - 16 years) of Karachi. Pak. j. rehabil. 2012;1(1):32-36]

- [4] Chunga MK, Leeb I, Keec D. Effect of stool height and holding time on postural load of squatting postures. *Int J IndErgon* 2003;32:309-317.
- [5] Hanvold TN, Veiersted KB, Wærsted M. A Prospective Study of Neck, Shoulder, and Upper Back Pain among Technical School Students Entering Working Life. *J Adolesc Health* 2010; 46:488-494.
- [6] Hurwitz EL, Aker PD, Adams AH, Meeker WC, Shekelle PG. Manipulation and mobilization of the cervical spine: a Sys Rev of the literature. *Spine* 1996;21(15):1746-59.
- [7] Benyamin R, Singh V, Parr AT, Conn A, Diwan S, Abdi S. Systematic Review of the Effectiveness of Cervical Epidurals in the Management of Chronic Neck Pain; *J Headache Pain* 2009;12:137-157.
- [8] Robbins M, Johnson IP, Cunliffe C. Encouraging good posture in school children using computers. *Chiropr Man Therap* 2009;12:35-44.
- [9] Grimmer-Somers K, Milanese S, Louw Quinette. Measurement of Cervical Posture in the Sagittal Plane. *J Manipulative Physiol Ther* 2008;31(7) 489-558
- [10] Selwyn N. The effect of using a home computer on students' educational use of IT. *J Med Internet Res* 1998;31(2):211-227
- [11] Gold JE, Driban JB, Yingling VR, Komaroff. a Characterization of posture and comfort in laptop users in non-desk settings, Elsevier Ltd and The Ergonomics Society 2011;43(2):392-9.
- [12] Posture and Study Habits Guide. McKinley Health Center University of Illinois. 2008. Retrieved from: http://www.mckinley.illinois.edu/handouts/pdfs/posture_study_habits.pdf
- [13] Grieco A. Sitting posture: an old problem and a new one. *Ergonomics* 1986;29:345-362.
- [14] Straker LM, Smith AJ, Bear N, O'Sullivan PB, de Klerk NH. Neck/shoulder pain, habitual spinal posture and computer use in adolescents: the importance of gender. *Ergonomics* 2011;54(6):539.
- [15] Parcells C, Hubbard MS. Mismatch of classroom furniture and student body dimensions: Empirical findings and health implications; *J Adolesc Health* 1999;24(4):265-273.
- [16] Smith L, Louw Q, Crous L, Grimmer-Somers K. Prevalence of neck pain and headaches: impact of computer use and other associative factors *Cephalalgia* 2009;29(2):250-257.
- [17] Edmondston SJ, Chan YH, Ngai GCW, Warren MLR, Williams JM, Netto SGK. Postural neck pain: An investigation of habitual sitting posture, perception of 'good' posture and cervicothoracic kinaesthesia; *Manual Therapy* 2007;12(4):363-371.
- [18] Cook C, Burgess-Limerick R. The effect of forearm support on musculoskeletal discomfort during call centre work. *Appl Ergon* 2004;35(4):337-42.
- [19] Chiung-Yu Cho, Survey of faulty postures and associated factors among Chinese adolescents *J Manipulative Physiol Ther*. 2008;31(3):224-9
- [20] Murphy S, Buckle P, Stubbs D. Classroom posture and self-reported back and neck pain in school children. *Appl Ergon*. 2004;35(2):113-20
- [21] Australian Bureau of Statistics. Children's participation in cultural and leisure activities, Australia. Report Number: 4901.0, Canberra, ACT: Australian Bureau of Statistics; 2012 <http://www.abs.gov.au/ausstats/abs@.nsf/mf/4901.0>
- [22] Balague F, Dutoit G, Waldburger M. Low back pain in school children: an epidemiological study. *Scandinavian J Rehabil Med* 1988;20:175-9.
- [23] Dockrell S, Earle D, Galvin R. Computer-related posture discomfort in primary school children: The effects of a school-based ergonomic intervention *J Med Internet Res* 2010;55:276-284
- [24] Hojat B and Mahdi E. Effect of different sitting posture on pulmonary function in students. *J Physiol* 2011;2(3):29
- [25] Murphy S, Buckle P, Stubbs D. Classroom posture and self-reported back and neck pain in school children. *App Ergonomics* 2004;35:113-120

INTRODUCTION

Worldwide childhood obesity is a considerable health problem^{1,2} and this epidemic is also becoming a challenging issue in developing countries like Pakistan³⁻⁵. WHO estimate that out of 1 billion overweight people in the world 300000 are obese^{6,7}. In the past a fat child was considered to be a healthy one but now obesity and over weight in children is major health concern because it may be responsible for many health problems in both childhood and adult hood. Obesity is associated with the onset of many diseases like diabetes mellitus, hypertension, dyslipidemia, polycystic ovarian disease, atherosclerosis^{1,8-10} and psychosocial problem^{8,11,12}. Many studies have shown that obesity may be responsible for the initiation of disease at an earlier stage^{6,13,14}. Another probable cause of obesity may be the genetic factor and life style¹⁵. A major problem which has been identified presently, is the conversion of active life style to sedentary life style, which itself is responsible in great majority for the incidence of obesity. Like outdoor games are replaced by computer and TV, junk food has become a fashion and a way of delight⁸.

A lot of work has been done on childhood obesity world-wide and in Pakistan. Recently, in 2012 a survey was conducted in Karachi children to determine the time given to physical activities in a day by these children the survey revealed that most of the children were spending only half an hour in physical activities among which the majority was boys¹⁶. According to the recommendation for the Asia pacific population B.M.I, greater than 23 will be considered as overweight whereas B.M.I, greater than 25 will be taken as obese rather than a usual B.M.I reference. In recent study using Indo-Asian specific cutoff values it is shown that quarter of Pakistan population would be counted as overweight and obese. Girls aged between 13-14 years spend most of their time in watching TV as compared to boys therefore are more prone to develop obesity¹⁶. In another study taking the B.M.I greater than 25 as obese¹⁴ the prevalence of obesity was found to be 18 percent in adolescence which coincides with the figure in the west⁶. National health survey of Pakistan 1990-94 and the Karachi survey 2004-05 reported prevalence of overweight in urban school children to be 3% and 5.7% respectively³. One of the major reason of overweight and obesity in school going children was due to decline in physical activity¹⁶.

Obesity is hard to reverse once established¹⁷⁻¹⁹. The risk of childhood overweight will continue into adolescence and adulthood. Having huge burden of obesity in the people in near future will be difficult to tackle. Timely preventive measures in children and adolescence will be sound and effective way of dealing with problem of obesity. School is the place where children spend half of the day therefore; teaching about the changes in lifestyle and including regular physical activity in the curriculum can be one of the benefits that can be reaped through this intervention. Family environment and parental characteristics such as parental working status sibling, person living in the child room etc must be considered in designing the strategies and intervention as it has strong influence on childhood obesity³. This study is focused to identify the various risk factors leading to overweight and obesity and to evaluate the role of Physical therapist in health education in schools.

MATERIAL AND METHODS

A cross sectional survey was conducted in school going school going children aged 13-16 years studying in

different schools of Karachi. A total of 300 students were selected through convenience sampling technique. Duration of study was one year. Data was collected by first taking permission from the head of the selected schools. Written Parental permission was taken from the selected students after explaining the objectives of the study. On the day of data collection all those students who were allowed by their parents to participate in the study and were present were assessed for height and weight measurement through calibrated weighing scale and height chart. BMI was calculated by using the standard formula of wt in kg/height in meter square. A self administered questionnaire in both English and Urdu was given to the students. The questionnaire assessed them about their routine activities, type of diet they are taking, number of meals/day, type of exercises they do, about physical activity in their schools and engagement in any sports activity. They were further asked about family history of obesity and their perceptions regarding the risk factors of obesity.

Data entry and analysis was done on SPSS version 20. Descriptive statistics were taken out for categorical variables in the form of percentages and frequencies.

RESULTS

A total of 300 participants were included in the study where number of girls and boys were almost equal (male=46.3 5% and female=53.7%). The children were incorporated from the schools in Karachi, aged between 13-16 years from different areas. At the time of recruitment a questioner was given to children for the assessment of factors leading to obesity. The height and weight of the individuals were taken to calculate the B.M.I. An awareness session for health and obesity was also incorporated and for that brochures were distributed among children.

BMI	Frequency	Percent
Underweight	67	22.3
Normal	121	40.3
Overweight	97	32.3
Obese	15	5.0
Total	300	100.0

It was found that majority 40% of the children come under the category of normal BMI while 32% were overweight, 22% were underweight and only 5% were found to be obese.

When asked about number of meals they take in a day 45% said 4 times, 34% affirmed 3 times a day and only 5.7% were eating 5 meals a day.

No. of time taking meal	Frequency	Percent
2 times	46	15.3
3 times	101	33.7
4 times	136	45.3
5 times	17	5.7
Total	300	100.0

On a question regarding the fast food consumption 40% children responded as occasionally, 30% stated rarely and 21.7% gave positive response for having fast food. Only 8% responded in negation.

Eating fast food	Frequency	Percent
none	23	7.7
rarely	90	30.0
occasionally	120	40.0
frequently	65	21.7
Total	300	100.0

It was observed that 48% children rarely took hawkers food, 24% avoided hawkers food while few stated occasionally (21.7) and frequently (6.7%).

Hawkers food	Frequency	Percent
none	72	24.0
rarely	143	47.7
occasionally	65	21.7
frequently	20	6.7
Total	300	100.0

Regarding daily water intake 42.3% children were drinking 6 glasses of water per day, 28% said more than 6 glasses per day while others consumed 4 or less than 4 glasses of water per day.

Daily water intake	Frequency	Percent
less than 4 glasses	7	2.3
4 glasses	81	27.0
6 glasses	127	42.3
more than 6 glasses	84	28.0
Total	300	100.0

When inquired about physical training period 53% of children have PT session, while other doesn't have physical therapy sessions in their time table.

physical training period in time table	Frequency	Percent
yes	159	53.0
no	136	45.3
Total	300	100.0

In response to TV watching habits, 36% claimed to spent more than 2 hrs, 25% spent more than 4 hours while others spent hour or more.

watching T.V in a day	Frequency	Percent
just 1 hr	50	16.7
more than 1 hr	67	22.3
more than 2 hr	107	35.7
more than 4 hr	76	25.3
Total	300	100.0

It was observed that 55% of the children have the positive family history of obesity, while 45% did not have obesity in their family.

Family history of obesity	Frequency	Percent
yes	165	55.0
no	135	45.0
Total	300	100.0

When enquired about perception of why children tend to get obese, 51% children regarded lack of physical activity as major cause of obesity while 48% considered improper diet. In others opinion family history (41%), improper sleep and medications were also the reasons for getting obesity.

Reasons of obesity in children prospective	Frequency	percentage
lacking physical activity	153	51%
having improper diet	145	48.3%
family history	123	41%
improper sleeping pattern	83	27.7%
taking medications	12	4%

On a question regarding physical activity participation 54% responded as yes, among which 43% were involved in different sports and 13% performed different exercises.

Engage in physical activity	Frequency	Percent
yes	163	54.3
no	137	45.7
Total	300	100.0

DISCUSSION

It is well know that obesity and overweight are associated with many risk factors such as improper dietary patterns, family history, physical inactivity and sedentary life style, lack of exercise, watching television or prolonged seated work etc. In the study we conducted, out of 300 children 97 were found to be overweight (32.3%) and 15 were obese (5%). In our study the frequency of overweight children was found to be high, 32.2%, when compared to the similar study conducted in Lahore, Pakistan which revealed 17%²⁰. The recent increase may be due to the increased commercials of fast foods and transformation

of active lifestyle to sedentary life style as in west. Even the studies conducted outside Pakistan shows lower incidence of overweight children as shown by the results of research conducted at Monastir where the overweight children ratio was 9.1% and in Tunis where the ratio was 19.7%^{21,22}. The frequency of obesity is however similar to the study in Tunis (5.7%)²².

Regarding junk food consumption our research showed that 40% children have it occasionally whereas a similar study conducted in Karachi in 2008 showed 80% children as junk food consumers²³. Our research also revealed that children lack the physical training and 45.3% schools didn't even had a physical training period as schedule. In addition, it was also observed that 50% of obese boys did not participate in physical activity while 63.6% girls also said that they do not take part in physical activities. A similar study advocated 81% obese boys and 78% obese girls do not affiliate themselves with physical activity²⁴. It is well understood that the major reason for putting on weight is high calorie intake and low output which is reflected in our society as obese and overweight children²⁴ and the reason is evident by the study that now a days we are having high calorie diets (junk food) but less physical workout indicating the lifestyle conversion into sedentary one. Sedentary pattern of life style promotes leisure activities which are unhealthy, such as using computer excessively, playing video games and watching different TV programmes. In return these shows get heavy advertisement adds of candies, snack food and sweetened breakfast cereals that are rich in calories and fats and lower in fiber and nutrients hence influencing children and promoting unhealthy diet pattern. It is also interesting that half of our study population has positive family history Hereditary contributes a major role as shown in several studies like in 2008 a study proclaimed for about 5-40% risk for obesity when parents were obese²⁴. Similarly another study indicated that there is 75% chance that a child will be overweight if both parents are obese and 20-50% chance if only single parent is obese. Another factor contributing to weight gain is sleeplessness; our study revealed that only 13.3% children had 8 or more than 8 hours sleep. This finding is also somewhat close to another study, conducted in the City of Trois-Rivières (Québec) in 2006²⁵ which stated 23.1% boy who sleeps 8-10 hours and 20.8% girls were obese and overweight.

CONCLUSION

Our study concluded that the major risk factor of obesity in 13-16 years children is lack of physical activity. It is also important to see that the usage of junk food is increasing in this age group that may be due to the enormous advertisement of junk stuffs. Another reason is the preference of relaxed/sedentary life style which is causing this problem in this population. Hereditary also plays an important part and our research also revealed that most of our population carries obesity in its genes. One very important finding is about the sleep pattern which has a strong relationship with weight gain.

REFERENCE

- [1] Gupta K, Shah P, Mishra A, Bharadwaj S et al. Secular trends in prevalence of overweight and obesity from 2006-2009 in urban Asian Indian adolescents aged 14-17 years. JPMC 2011;doi:10.1371/journal.pone.0017221. [available online] <http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0017221>
- [2] Frayling MT, Nicholas J, Timpson, NJ Weedon MM. A

- A Common Variant in the FTO Gene Is Associated with Body Mass Index and Predisposes to Childhood and Adult Obesity 2007;316 : 889-894.
- [3] Mushtaq MU, Gull S, Abdullah HM, Shahid U, Shad MA, Akram J. Prevalence and socioeconomic correlates of overweight and obesity among Pakistani primary school children. BMC Public Health 2011; 11:724 doi:10.1186/1471-2458/11/724
- [4] Qidwai W, Ashfaq T. Imminent Epidemic of Diabetes Mellitus in Pakistan: Issues and Challenges for Health Care Providers, JLUHMS 2010;9(3):231-34.
- [5] Mickey Chopra, Sarah Galbraith & Ian Darnton-hill A global response to a global problem: the epidemic of overnutrition. Bulletin of the World Health Organization;2002.34
- [6] Warraich HJ, Javed F, Faraz-ul-Haq M, Khawaja FB, Saleem S. Prevalence of Obesity in School-Going Children of Karachi. PLoS ONE 2009;4(3):e4816. doi:10.1371/journal.pone.0004816.
- [7] Puska P, obesity and overweight. Global strategy on diet, physical activity and health 2003. Volume 376, Issue 9754, 20-26 November 2010, Pages 1775-1784
- [8] Hoeger WK & Hoeger A principal and labs for fitness and wellness. 6th Ed. USA: wadsworth; 2010.202-05.
- [9] Levy TS, Raun CM, Castellanos CA, Coronel AS, Aguilar AJ, Humaranl MG. Effectiveness of a diet and physical activity promotion strategy on the prevention of obesity in Mexican school children. BMC public health 2012;12:152 doi:10.1186/1471-2458/12/152.
- [10] Halpern A, Mancini CM, Magalhães CE. Metabolic syndrome, dyslipidemia, hypertension and type 2 diabetes in youth: from diagnosis to treatment. licensee BioMed Central Ltd 2010;2 2:55 doi:10.1186/1758-5996-2-55
- [11] Aronne LJ, Segal KR. Weight Gain in the Treatment of Mood Disorders. Journal of Clinical Psychiatry 2003; 64(8):22-29.
- [12] Bell SJ, Goodrick GK. A Functional Food Product for the Management of Weight. Critical Reviews in Food Science and Nutrition 2002;42:163-178
- [13] Wahba M, Mak H. Obesity and Obesity-Initiated Metabolic Syndrome: Mechanistic Links to Chronic Kidney Disease. clinical journal of american society of nephrology 2007;2:550-562
- [14] American Obesity Association. 2002. What is obesity? Retrieved September 25, 2006, from http://www.obesity.org/information/what_is_obesity.asp
- [15] Frayling MT, Nicholas J, Timpson NJ, Weedon MM. A Common Variant in the FTO Gene Is Associated with Body Mass Index and Predisposes to Childhood and Adult Obesity 2007;316 : 889-894.
- [16] Jafar TH, Qadr ZI, Islam M, Hatcher J, Bhutta ZA, Chaturvedi N. Rise in childhood obesity with persistently high rates of under nutrition among urban school-aged Indo-Asian children. Arch Dis Child 2008;93:373-378 doi:10.1136/adc.2007.125641. Available from :<http://adc.bmj.com/content/93/5/373.full>
- [17] Pletzer K, Pengpid S. Overweight and obesity and associated factors among school aged adolescents in Ghana and Uganda. Inj Environ Res public health 2011;8,3859-3870 doi:10.3390/ijerph8103859.
- [18] Peltzer K, Pengpid S. [homepage on the Internet]. 2011 cited 2012 Jul 2. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3210586/>
- [19] Waters E. Interventions for preventing obesity in children Review. The Cochrane Library 2011;(12):1-70.
- [20] Chaput J-P, Brunet M and Tremblay A. Relationship

between short sleeping hours and childhood overweight obesity results from, the quebec form project. International journal of obesity 2009;30 Doi: 10.1038/sj.ijo.0803291.

- [21] Abdelkafi KA, Younes K, Gabzi Z, Bouslah A, Maalell, Maatouk EIMW, Dahmen H, Bel AN, Gabzi A, et al. Risk factor of children overweight and obesity. Tunis Med 2012 May;90(5):387-93.
- [22] Boukthir S, Essaddam L, MazighMrad S, Ben Hassine L, Gannouni S, Nessib F, Bouaziz A, Brini I, Sammoud A, Bouyahia O, Zouari B. Prevalence and risk factors of overweight and obesity in elementary school children in the metropolitan region of Tunis, Tunisia. Tunis Med 2011,Jan;89(1):50-4

- [23] Warraich HJ, Javed F, Faraz-ul-Haq M, Khawaja FB, Saleem S Prevalence of Obesity in School-Going Children of Karachi. PLoS ONE 2009;4(3):e4816. doi:10.1371/journal.pone.0004816.
- [24] Childhood Obesity Assistant Secretary for Planning and Evaluation ASPE.hhs.gov US department of Health & Human Services. 2005 [available online]: http://aspe.hhs.gov/health/reports/child_obesity/
- [25] Childhood obesity by Mayo clinic staff. May 4 2012 available online at: <http://www.mayoclinic.com/health/childhood-obesity/DS00698/DSECTION=risk-factors>